ERRATA

1. Ecological Restoration 30:1 (March 2012): In Snee et al.'s Restoration Note entitled, "Passive Soil Manipulation Influences the Successional Trajectories of Forest Communities at a Denuded Former Campsite," the following reference was omitted:

Ewing, K. 2002. Effects of initial site treatments on early growth and three-year survival of Idaho fescue. *Restoration Ecology* 10:282–288.

2. Ecological Restoration 30:2 (June 2012). In Moser et al.'s article entitled, "Control of Hardwood Regeneration in Restored Carolina Bay Depression Wetlands," the full list of references and the Appendix were omitted. We include them here:

Appendix

Identified Species Lists Before and After Restoration Before Restoration Species List†

Common name(s)	Scientific name	Wetland indicator status *
red maple	Acer rubrum	FAC
peppervine	Ampelopsis arborea	FAC+
American beautyberry	Callicarpa americana	FACU-
common persimmon	Diospyros virginiana	FAC
yellow jessamine	Gelsemium sempervirens	FAC
American holly	llex opaca var. opaca	FAC-
sweetgum	Liquidambar styraciflua	FAC+
partridgeberry	Mitchella repens	FACU+
wax myrtle	Myrica cerifera	FAC+
swamp tupelo	Nyssa sylvatica var. biflora	OBL
maidencane	Panicum hemitomon	OBL
loblolly pine	Pinus taeda	FAC
laurel oak	Quercus laurifolia	FACW
water oak	Quercus nigra	FAC
willow oak	Quercus phellos	FACW-
catbrier	Smilax pumila	N.I.‡
roundleaf greenbrier	Smilax rotundifolia	FAC
Elliott's blueberry	Vaccinium elliottii	FAC+
muscadine	Vitis rotundifolia	FAC

[†]There was also 1 Carex sp. which could not be identified past genus level.

After Restoration Species List†

Common name	Scientific name	Wetland indicator status *
red maple	Acer rubrum	FAC
annual ragweed	Ambrosia artemisiifolia	FACU
peppervine	Ampelopsis arborea	FAC+
greenwhite sedge	Carex albolutescens	FAC+
southern waxy sedge	Carex glaucescens	OBL
slender woodoats	Chasmanthium laxum	FACW-
Elliott's croton	Croton elliottii	FACW+
Baldwin's flatsedge	Cyperus croceus	FAC
redroot flatsedge	Cyperus erythrorhizos	OBL
strawcolored flatsedge	Cyperus strigosus	FACW
openflower rosette grass	Dichanthelium laxiflorum	FAC
velvet panicgrass	Dichanthelium scoparium	FACW

^{*-} Region 2 wetland indicator status

[‡]N.I.- no indicator status available

Common name	Scientific name	Wetland indicator status *
roughhair rosette grass	Dichanthelium strigosum var. leucoblepharis	FAC
common persimmon	Diospyros virginiana	FAC
fireweed	Erechtites hieraciifolia	FAC-
dog-fennel	Eupatorium capillifolium	FACU
green ash	Fraxinus pensylvanica	FACW
yellow jessamine	Gelsemium sempervirens	FAC
American pennywort	Hydrocotyle americana	OBL
dwarf St. John's wort	Hypericum mutilum	FACW
American holly	llex opaca var. opaca	FAC-
needlepod rush	Juncus scirpoides	FACW+
grassleaf rush	Juncus marginatus	FACW
sweetgum	Liquidambar styraciflua	FAC+
Japanese honeysuckle	Lonicera japonica	FAC-
seedbox	Ludwigia alternifolia	OBL
marsh seedbox	Ludwigia palustris	OBL
axilflower	Mecardonia acuminata	FACW
wax myrtle	Myrica cerifera	FAC+
swamp tupelo	Nyssa sylvatica var. biflora	OBL
Bosc's mille graines	Oldenlandia boscii	FACW-
clustered mille graines	Oldenlandia uniflora	FACW-
warty panicgrass	Panicum verrucosum	FACW
Virginia creeper	Parthenocissus quinquefolia	FAC
American Pokeweed	Phytolacca americana	FACU+
loblolly pine	Pinus taeda	FAC
water elm	Planera aquatica	OBL
swamp smartweed	Polygonum hydropiperoides	OBL
bald-rush	Psilocarya nitens	OBL
laurel oak	Quercus laurifolia	FACW
water oak	Quercus nigra	FAC
willow oak	Quercus phellos	FACW-
Maryland meadowbeauty	Rhexia mariana	FACW+
handsome Harry	Rhexia virginica	FACW+
winged sumac	Rhus copallinum	N.I.‡
anglestem beaksedge	Rhynchospora caduca	OBL
nodding beaksedge	Rhynchospora inexpansa	FACW
pineland beaksedge	Rhynchospora perplexa	OBL
sand blackberry	Rubus cuneifolius	FACU
fringed nutrush	Scleria ciliata	FAC
reticulated nutrush	Scleria reticularis	OBL
fewflower nutrush	Scleria pauciflora	FAC+
sassafras	Sassafras albidum	FACU
catbrier	Smilax pumila	N.I.
roundleaf greenbrier	Smilax rotundifolia	FAC
poison oak	Toxicodendron pubescens	FACU
Elliott's blueberry	Vaccinium elliottii	FAC+
	Viola lanceolata	OBL
bog white violet muscadine	Viola lanceolata Vitis rotundifolia	FAC
muscadine	vitis roturialiolia	FAC

[†]There were also 1 Eleocharis sp., 1 Hypericum sp., 1 Isoetes sp., 3 Juncus spp., 1 Rhexia sp., and 2 Rhynchospora spp. that could not be identified past genus level. 10 sterile Gramineae family specimens were also collected that were left at the family level of taxonomic identification (several of these 10 specimens are suspected to be the same species). *- Region 2 wetland indicator status

[‡]N.I.- no indicator status available

References

- Barton, C.D., D.D. DeSteven and J.C. Kilgo. 2004. Mitigation bank promotes research on restoring coastal plain depression wetlands (South Carolina). *Ecological Restoration* 22: 291–292.
- Barton, C.D., D.M. Andrews and R.K. Kolka. 2008. Evaluating hydroperiod response in restored Carolina bay wetlands using soil physiochemical properties. *Restoration Ecology* 16: 668–677.
- Barton, C.D., D.D. De Steven, R. Sharitz, J. Kilgo, K. Kinkead, D. Otis, H. Hanlin, J. Ledvina, B. Taylor and J. Blake. 2007. The Carolina Bay Restoration Project: Final Report 2000–2006. US Department of Energy DE-A109-00SR22188US. Department of Energy's Information Bridge: www.osti.gov/bridge/product .biblio.jsp?query_id=2&page=0&osti_id=921084.
- Barton, C.D., J.J. Blake and D. W. Imm. 2005. Ecological restoration. Pages 84–102 in J.C. Kilgo and J.I. Blake, (eds), *Ecology and Management of a Forested Landscape: Fifty Years on the Savannah River Site* Washington DC: Island Press.
- Battaglia, L.L. and B.S. Collins. 2006. Linking hydroperiod and vegetation response in Carolina bay wetlands. *Plant Ecology* 184: 173–185.
- BASF Corporation. 2005. HABITAT® herbicide for aquatic and invasive vegetation control: Technical bulletin. Research Triangle Park, NC: BASF Corporation.
- Beavers, J.R., J.G. Hensley, J.G. Byrne, E.E. Clebsch and J.H. Jenkins. 1977. Opportunities for resource management - an ecological analysis of the Savannah River Plant. Atlanta, Georgia: Southern Regional Office, United States Forest Service.
- Bennett, S.H. and J.B. Nelson. 1991.
 Distribution and status of Carolina bays in SouthCarolina. Cloumbia, SC: Nongame and Heritage Trust Publications, South Carolina Wildlife and Marine Resources Department.
- Blake, J.I., C H. Hunter and B A. Bayle. 2005. Climate and air quality. Pages 20–30 in J.C. Kilgo and J.I. Blake, (eds), Ecology and Management of a Forested Landscape: Fifty Years on the Savannah River Site. Washington DC: Island Press.
- Bliss, C.M. and N.B. Comerford. 2002. Forest harvesting influence on water

- table dynamics in a Florida flatwoods landscape. *Soil Science Society of America Journal* 66: 1344–1349.
- Burns, R.M. and B.H. Honkala. 1990.
 Silvics of North America: 1. Conifers;
 2. Hardwoods. Agriculture Handbook
 654. Washington DC: U.S.
 Department of Agriculture, Forest
 Service.
- Collins, B.S. and L.L. Battaglia. 2001. Hydrology effects on propagule bank expression and vegetation in six Carolina bays. *Community Ecology* 2: 21–33.
- Dietz-Brantley, S.E., B.E. Taylor, D.P. Batzer and A.E. DeBiase. 2002. Invertebrates that aestivate in dry basins of Carolina bay wetlands. *Wetlands* 22: 767–775.
- De Steven, D., R.R. Sharitz and C.D. Barton. 2010. Ecological outcomes and evaluation of success in passively restored southeastern depressional wetlands. *Wetlands* 30: 1129–1140.
- De Steven, D.D. and R.R. Sharitz. 2007. Transplanting native dominant plants to facilitate community development in restored Coastal Plain wetlands. *Wetlands* 27: 972–978.
- De Steven, D., R.R. Sharitz, J.H. Singer and C.D. Barton. 2006. Testing a passive revegetation approach for restoring coastal plain depression wetlands. *Restoration Ecology* 14: 452–460.
- De Steven, D. and M.M. Toner. 2004. Vegetation of upper coastal plain depression wetlands: Environmental templates and wetland dynamics within a landscape framework. Wetlands 24: 23–42.
- Drew, M.B., P.C. Goebel, L.K. Kirkman, B.J. Palik and L. West. 2000.

 Depressional wetland vegetation types: A question of plant community development. *Wetlands* 20: 373–385.
- Fargione, J. and D. Tilman. 2002.
 Competition and coexistence in terrestrial plants. Pages 165–206 in U. Sommer and B. Worm. (eds) Ecological Studies, vol. 161.
 Competition and Coexistence. Berlin Heidelberg: Springer-Verlag.
- Gibbons, J.W. 2003. Terrestrial habitat: A vital component for herpetofauna of isolated wetlands. *Wetlands* 23: 630–635.
- Hook, D.D. and J.R. Scholtens. 1978.
 Adaptations and flood tolerance of tree species. Pages 299–331 in D.D.
 Hook and R.M.M. Crawford (eds)
 Plant Life in Anaerobic Environments.

- Ann Arbor, MI: Ann Arbor Science Publishers.
- Imm, D.W. and K.W. McLeod. 2005.
 Plant communities. Pages 106–161
 in J.C. Kilgo and J.I. Blake, (eds),
 Ecology and Management of a Forested
 Landscape: Fifty Years on the Savannah
 River Site Washington DC: Island
 Press.
- Kilgo, J.C. and J.I. Blake. 2005. Ecology and Management of a Forested Landscape: Fifty Years on the Savannah River Site Washington DC: Island Press.
- Kirkman, L.K. 1995. Impacts of fire and hydrological regimes on vegetation in depression wetlands of southeastern USA. Pages 10–20 in S.I. Cerulean and R.T. Engstrom (eds) Fire in Wetlands: A Management Perspective. Proceedings of the Tall Timbers Fire Ecology Conference, No. 19. Tallahassee, FL: Tall Timbers Research Station.
- Kirkman, L.K., R.F. Lide, G.R. Wein and R.R. Sharitz. 1996. Vegetation changes and land-use legacies of depression wetlands of the western coastal plain of South Carolina: 1951–1992. Wetlands 16: 564–576.
- Kirkman, L.K and R.R. Sharitz. 1994. Vegetation disturbance and maintenance of diversity in intermittently flooded Carolina bays in South Carolina. *Ecological Applications* 4: 177–188.
- Lide, R.F. 1997. When is a depressional wetland a Carolina bay? *Southeastern Geography* 37: 90–98.
- McLeod, K.W., M.R. Reed and L.D. Wilke. 2000. Elevation, competition control, and species affect bottomland forest restoration. *Wetlands* 20: 162–168.
- Passarge, J. and J. Huisman. 2002.
 Competition in well-mixed habitats:
 From competitive exclusion to competitive chaos. Pages 7–42
 in U. Sommer and B. Worm
 (eds) Ecological Studies, Vol. 161.
 Competition and Coexistence Berlin Heidelberg: Springer-Verlag.
- Poiani, K.A. and P.M. Dixon. 1995. Seed banks of Carolina bays: Potential contributions from surrounding landscape vegetation. *American Midland Naturalist* 134: 140–154.
- Pyzoha, J.E., T.J. Callahan, G. Sun, C.C. Trettin and M. Miwa. 2008. A conceptual hydrologic model for a forested Carolina bay depressional wetland on the coastal plain of South

- Carolina, USA. Hydrological Processes 22:2689-2698.
- Schalles, J.F., R.R. Sharitz, J.W. Gibbons, G.J. Leversee and J.N. Knox. 1989. Carolina Bays of the Savannah River Plant. SRO-NERP-18 Aiken, SC: Savannah River Ecology Laboratory.
- Schalles, J.F. and D.J. Shure. 1989. Community structure and productivity patterns of a dystrophic Carolina bay wetland. Ecological Monographs 59: 365-385.
- Sharitz, R.R. 2003. Carolina bay wetlands: Unique habitats of the southeastern United States. Wetlands 23: 550-562.
- Snodgrass, J.W., M.J. Komoroski, A.L. Bryan Jr. and J. Burger. 2000. Relationships among isolated wetland size, hydroperiod, and amphibian species richness: Implications for wetland regulations. Conservation Biology 14: 414-419.
- Sullivan, T.P. and D.S. Sullivan. 2003. Vegetation management and

- ecosystem disturbance: Impact of glyphosate herbicide on plant and animal diversity in terrestrial systems. Environmental Reviews 11: 37-59
- Sun, G.C., T.J. Callahan, J.E. Pyzoha and C.C. Trettin. 2006. Modeling the climatic and subsurface stratigraphy controls on the hydrology of a Carolina bay wetland in South Carolina, USA. Wetlands 26: 567-580.
- Sun, G.C., H. Riekerk and N. B. Comerford. 1998. Modeling the hydrologic impacts of forest harvesting on Florida flatwoods. Journal of the American Water Resources Association 34: 843–854.
- Sun, G.C., H. Riekerk and L.V. Kornhak. 2000. Ground-water-table rise after forest harvesting on cypress-pine flatwoods in Florida. Wetlands 20: 101-112.
- Taylor, B.E., D.A. Leeper, M.A. McClure and A.E. DeBiase. 1999. Carolina

- bays: Ecology of aquatic invertebrates and perspectives on conservation. Pages 167-196 in R.B. Rader, D.P. Batzer and S.A. Wissinger (eds) Invertebrates in Freshwater Wetlands of North America: Ecology and Management. New York, NY: John Wiley and Sons, Inc.
- Tiner, R.W. 2003. Geographically isolated wetlands of the United States. Wetlands 23: 494-516.
- Williams, T.M. and D.J. Lipscomb. 1981. Water table rise after cutting on coastal plain soils. Southern Journal of Applied Forestry 5: 46-48.
- Xu, Y.J., J.A. Burger, W.M. Aust, S.C. Patterson, M. Miwa and D.P. Preston. 2002. Changes in surface water table depth and soil physical properties after harvest and establishment of loblolly pine (Pinus taeda L.) in Atlantic coastal plain wetlands of South Carolina. Soil and Tillage Research 63: 109-121.